



OPV: oral poliovirus  
 DTaP: diphtheria and tetanus toxoids and acellular pertussis vaccine  
 DT: diphtheria and tetanus toxoids  
 BCG: bacille Calmette-Guérin  
 PPD: purified protein derivative

Schedule has been changed due to the revision of Tuberculosis Prevention Law in April 2005.

↓ shot   ■ age regularly vaccinated   □ age vaccination is regulated   □ (dotted) age vaccinated   □ (dotted) program of prevention of mother-to-child infection

\*1 Category I: The objective is to protect the public health. Immunization is strongly recommended by law.  
 \*2 Category II: The objective is to protect the individual's health. This is available on request.  
 \*3 60-64 year-old persons who have severe disorders of the heart, kidney, respiratory organs, or from the human immunodeficiency virus

\*4 Newborn babies from HBs antigen positive mothers should receive anti-HBs human immunoglobulin (HBIG) and HB vaccine. HBIG is intramuscularly injected to the neonates within 48 hr and 2-3 months after birth, followed by hepatitis B vaccine starting at the time of the second immunoglobulin injection. The vaccine is given three times; the first two doses at one month apart and the third one in three months after the first dose. This is covered by health insurance.  
 © Copyright 2005 IDSC All Rights Reserved.